

## CLAIMS

1. A method of identifying the presence of CD94/NKG2+ NK  
5 cells and T cells in a sample, which method comprises contacting the  
sample with HLA-E under binding conditions and detecting binding of HLA-  
E to the cells.
2. The method according to claim 1, wherein the HLA-E is  
labeled with a signal moiety and bound HLA-E is detected
- 10 3. A method of selecting CD94/NKG2+ cells from a sample,  
which method comprises contacting the sample with HLA-E under binding  
conditions and separating cells bound to the HLA-E from the sample.
4. The method according to claim 3, wherein the HLA-E is  
immobilised on a support.
- 15 5. A method of killing or inactivating CD94/NKG2+ cells, which  
method comprises contacting the cells with HLA-E under binding  
conditions and carrying out targeted killing on the bound cells.
6. The method according to claim 5, wherein the HLA-E is  
attached to an effector agent.
- 20 7. The method according to claim 6, wherein the effector agent  
is a toxic moiety which kills or inactivates the bound cells.
8. The method according to <sup>claim 5</sup> ~~any one of claims 5 to 7~~, wherein  
the CD94/NKG2+ cells are provided in a mixed cell population and the  
non-CD94/NKG2+ cells are recovered.
- 25 9. A method of modifying NK cell activity against a potential  
target cell, which method comprises expressing HLA-E at the target cell  
surface.
10. The method according to claim 9, wherein the HLA-E is  
expressed by a heterologous DNA stably integrated into the cell.

11. The method according to claim 10, wherein the cell does not normally express HLA-E.

12. The method according to claim 11, wherein the cell is a non-human mammalian cell.

13. The method according to <sup>claim 9</sup> ~~any one of claims 9 to 12~~, wherein the cell is present in a group of cells in a xenogeneic organ or tissue.

14. CD94/NKG2+ cells isolated by the method according to claim 3 ~~or claim 4~~.

15. A population of cells depleted of CD94/NKG2+ cells by the method according to claim 8.

16. A therapeutic method comprising introducing an effective quantity of cells according to claim 14 ~~or claim 15~~ into a patient.

17. A method comprising removing a sample of cells from a patient, isolating from the sample a population of CD94/NKG2+ cells or a population of CD94/NKG2 depleted cells, by the method according to claim 3 ~~or claim 4, or claim 8~~, and reintroducing the isolated population of cells into the patient.

18. A non-human mammalian cell which expresses HLA-E at the cell surface by virtue of a nucleic acid encoding HLA-E integrated into the genome of the cell.

19. A non-human mammal comprising cells according to claim 18.

20. A method of testing a compound for biological activity, which method comprises:

- 25
- (i) providing cells expressing CD94/NKG2 receptors at the cell surface;
  - (ii) contacting the cells with HLA-E in the presence of the test compound; and
  - (iii) determining whether the presence of the compound affects the binding of HLA-E to the cells.
- 30

21. The method according to claim 20, wherein the CD94/NKG2 receptors are inhibitory NK cell receptors such as CD94/NKG2A receptors.

22. The method according to claim 20, wherein the CD94/NKG2A receptors are stimulatory NK cell receptors such as CD94/NKG2C receptors.

23. Compounds identified by the method according to ~~any one of~~ <sup>claim 20</sup>

~~claims 20 to 22~~, as affecting the binding of HLA-E to CD94/NKG2 receptors. <sup>research</sup>

24. A multimer of HLA-E comprising two or more HLA-E molecules, said multimer having enhanced binding capability compared to non-multimeric HLA-E, optionally labelled with a signal moiety.

25. The multimer according to claim 24, comprising recombinant HLA-E molecules attached via a linker molecule.

26. The multimer according to claim 25, wherein the HLA-E molecules are biotinylated and attached via a linker molecule such as streptavidin, avidin or extravidin.

27. HLA-E coupled to a toxic agent.

28. The HLA-E according to claim 27, wherein the HLA-E is in the form of a multimer of HLA-E according to ~~any one of claims 24 to 26~~ <sup>claim 24</sup>

29. HLA-E immobilised on a support.